



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Towhee. On March 22, my friend, Mr. John De Q. Briggs, saw two Towhees at Plymouth, Mass. As they do not usually arrive in Massachusetts before the 20th of April, it is probable that they had wintered in that region. — ARTHUR C. COMEY, *Cambridge, Mass.*

Colorado Bird Notes.—I desire to record a Western Blue Grosbeak (*Guiraca caerulea eurhyncha*) taken near Altona, Boulder Co., Colo., August 16, 1901. The farthest north this bird had previously been observed in Colorado was at Morrison.

I wish also to record the observation of an Indigo Bunting (*Cyanospiza cyanea*) near Clear Creek, Denver, Colo., May 7, 1901. This is the fourth record for Colorado.

May 19, 1900, I found twelve Forster's Terns (*Sterna forsteri*) breeding at Barr, Colo. Mr. Ridgway states (Bull. Essex Institute, V. Nov. 1873, 174) that a few were found breeding in the State. I have found no records of their breeding here since that time.

Also at Barr, on June 20, 1900, I found a set of Canvasback's (*Aythya vallisneria*) eggs, and on July 4 I found another set, which was apparently laid by the same bird. I was informed upon good authority that there were two other pairs breeding in the vicinity. As far as I can determine this is the first record of the Canvasback's breeding in Colorado.—A. H. FELGER, *Denver, Colo.*

Some Southern New Hampshire and Western Massachusetts Notes.—The young bird student who has developed comparatively good observing powers, but has as yet no reputation, is unfortunate if he is made sole witness to interesting bird happenings which cannot be authenticated. I cannot hope that the following will all be accepted as records; for, by singularly bad luck, the bird was not secured in any one of the more interesting cases; and I can only wish they had fallen to the lot of some trusted man.

NEW HAMPSHIRE.

On July 28, 1900, I saw on the shore of a small lake (Nubanusit Lake) in Hillsboro County, southwestern New Hampshire, just over the line from Cheshire County, a Lincoln's Sparrow (*Melospiza lincolni*), whose actions seem to prove it a breeding bird.

Nubanusit Lake, partly in Cheshire and partly in Hillsboro County, is a deepish sheet of clear water, of irregular form, being nearly three miles long and varying in width from less than a tenth of a mile to about a mile and a quarter. It lies at a height above the sea of 1368 feet, while some of the spruce-clad hills by which it is surrounded reach a height of nearly 2300 feet.

The lake shore, which is now almost entirely wild, is here and there swampy and bush-grown, but mainly covered by a dense forest of fair-